

Weekly Exercises 0

Room: HF-115

Wednesday, 26.04.2017, 12:15-14:00

Reminder: Finite differences

Remember the Taylor approximation: Let $f \in C^{n+1}(\mathbb{R})$ be a real valued function that is $n+1$ -times continuously differentiable. Then there exists a function h_n such that

$$f(x) = \sum_{k=0}^n \frac{(x-y)^k}{k!} f^{(k)}(y) + \frac{(x-y)^{n+1}}{(n+1)!} f^{(n+1)}(\xi)$$

for some ξ between x and y .

Let f be two times differentiable, and let the second derivative be bounded by C , i.e. $\sup_x |f''(x)| \leq C$. Show that for any pair x_k, x_{k+1} with $h = x_{k+1} - x_k$ it holds that

$$f'(x_k) = \frac{f(x_{k+1}) - f(x_k)}{h} + r(h),$$

for some function r . Furthermore, it holds that $r(h) \in \mathcal{O}(h)$, i.e. there exists a constant c such that $|r(h)| \leq c|h|$ (for all sufficiently small h).

Intro to Sparse Linear Operators in MATLAB

Throughout the course we will work in the finite dimensional setting, i.e. we discretely represent gray value images $f : \Omega \rightarrow \mathbb{R}$ or color images $f : \Omega \rightarrow \mathbb{R}^3$ as (vectorized) matrices $f \in \mathbb{R}^{m \times n}$ ($\text{vec}(f) \in \mathbb{R}^{mn}$) respectively $f \in \mathbb{R}^{m \times n \times 3}$ ($\text{vec}(f) \in \mathbb{R}^{3mn}$). To discretely express functionals like the total variation for smooth f

$$TV(f) := \int_{\Omega} \|\nabla f(x)\| \, dx$$

you will therefore need a discrete gradient operator

$$D := \begin{pmatrix} D_x \\ D_y \end{pmatrix}$$

for vectorized representations $\text{vec}(f)$ of images $f \in \mathbb{R}^{m \times n}$ so that

$$TV(f) = \|D \text{vec}(f)\|_{2,1} = \sum_{i=1}^{nm} \sqrt{(D_x \cdot \text{vec}(f))_i^2 + (D_y \cdot \text{vec}(f))_i^2}.$$

The aim of this exercise is to derive the gradient operator and learn how to implement it with MATLAB.

Exercise 1. Let $f \in \mathbb{R}^{m \times n}$ be a discrete grayvalue image. Your task is to find matrices \tilde{D}_x and \tilde{D}_y for computing the forward differences f_x, f_y in x and y -direction of the image f with Neumann boundary conditions so that:

$$f_x = f \cdot \tilde{D}_x := \begin{pmatrix} f_{12} - f_{11} & f_{13} - f_{12} & \cdots & f_{1n} - f_{1(n-1)} & 0 \\ f_{22} - f_{21} & \cdots & & & 0 \\ \vdots & & & \vdots & 0 \\ f_{m2} - f_{m1} & \cdots & & f_{mn} - f_{m(n-1)} & 0 \end{pmatrix} \quad (1)$$

and

$$f_y = \tilde{D}_y \cdot f = \begin{pmatrix} f_{21} - f_{11} & f_{22} - f_{12} & \cdots & f_{2n} - f_{1n} \\ f_{31} - f_{21} & \cdots & & f_{3n} - f_{2n} \\ \vdots & & & \vdots \\ f_{m1} - f_{(m-1)1} & \cdots & & f_{mn} - f_{(m-1)n} \\ 0 & \cdots & 0 & 0 \end{pmatrix}. \quad (2)$$

Exercise 2. Implement the derivative operators from the previous exercise using MATLABs `spdiags` command. Load the image from the file `Vegetation-028.jpg` using the command `imread` and convert it to a grayvalue image using the command `rgb2gray`. Finally apply the operators to the image and display your results using `imshow`.

For our algorithms it is more convenient to represent an image f as a vector $\text{vec}(f) \in \mathbb{R}^{mn}$, that means that the columns of f are stacked one over the other.

Exercise 3. Derive a gradient operator

$$D = \begin{pmatrix} D_x \\ D_y \end{pmatrix}$$

for vectorized images so that

$$D_x \cdot \text{vec}(f) = \text{vec}(f_x) \quad D_y \cdot \text{vec}(f) = \text{vec}(f_y)$$

You can use that it holds that for matrices A, X, B

$$AXB = C \iff (B^\top \otimes A)\text{vec}(X) = \text{vec}(C)$$

where \otimes denote the Kronecker (MATLAB: `kron`) product.

Experimentally verify that the results of Ex. 2 and Ex. 3 are equal by reshaping them to the same size using MATLABs `reshape` or the `:` operator, and showing that the norm of the difference of both results is zero.

Exercise 4. Assemble an operator D^c for computing the gradient (or more precisely the Jacobian) of a color image $f \in \mathbb{R}^{n \times m \times 3}$ using MATLABs `cat` and `kron` commands.

Exercise 5. Compute the color total variation given as

$$TV(f) = \|D^c \text{vec}(f)\|_{F,1} = \sum_{i=1}^{nm} \left\| \begin{pmatrix} (D_x \cdot \text{vec}(f_r))_i & (D_x \cdot \text{vec}(f_g))_i & (D_x \cdot \text{vec}(f_b))_i \\ (D_y \cdot \text{vec}(f_r))_i & (D_y \cdot \text{vec}(f_g))_i & (D_y \cdot \text{vec}(f_b))_i \end{pmatrix} \right\|_F$$

of the two images `Vegetation-028.jpg` and `Vegetation-043.jpg` and compare the values. What do you observe? Why?